

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A23L 1/305, A61K 31/195, A23K 1/16	A1	(11) International Publication Number: WO 98/06278 (43) International Publication Date: 19 February 1998 (19.02.98)
(21) International Application Number: PCT/IB97/01117 (22) International Filing Date: 12 August 1997 (12.08.97) (30) Priority Data: 9616910.7 12 August 1996 (12.08.96) GB 9621914.2 21 October 1996 (21.10.96) GB (71) Applicant: HARRIS, Roger [GB/GB]; 4 Armstrong Close, Newmarket, Suffolk CB8 8HD (GB). (71)(72) Applicants and Inventors: DUNNETT, Mark [GB/GB]; 78 Highwood Road, Gazeley, Newmarket, Suffolk CB8 8RJ (GB). JOHANSSON, Kenny [SE/SE]; Fairing Sport Nutrition, ST Eriksgatan 16, S-112 39 Stockholm (SE). (74) Agent: DOST, Wolfgang; Bardehle Pagenberg Dost Altenburg Frohwitter Geissler, Postfach 86 06 20, D-81663 München (DE).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHODS AND COMPOSITIONS FOR INCREASING THE ANAEROBIC WORKING CAPACITY IN TISSUES (57) Abstract A method for increasing the synthesis and accumulation of beta-alanylhistidine dipeptides, with a simultaneous increase in the accumulation of creatine, in bodily tissues of humans and animals is described. This is accomplished by causing an increase in the blood plasma concentrations of beta-alanine and creatine, or the blood plasma concentrations of beta-alanine, L-histidine and creatine, by the ingestion or infusion of a composition including beta-alanine, beta-alanine and creatine, or beta-alanine, L-histidine and creatine, or active derivatives thereof.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHODS AND COMPOSITIONS FOR INCREASING THE ANAEROBIC
WORKING CAPACITY IN TISSUES

5 Background of the Invention

The invention relates to methods and compositions for increasing the anaerobic working capacity of muscle and other tissues.

10 Natural food supplements are typically designed to compensate for reduced levels of nutrients in the modern human and animal diet. In particular, useful supplements increase the function of tissues when consumed. It can be particularly important to supplement the diets of particular classes of animals whose the normal diet may be deficient in
15 nutrients available only from meat and animal produce (e.g., human vegetarians and other animals consume an herbivorous diet).

 For example, in the sporting and athletic community, natural food supplements which specifically improve athletic
20 ability are increasingly important, such as supplements that promote or enhance physical prowess for leisure or employment purposes. In another example, anaerobic (e.g., lactate-producing) stress can cause the onset of fatigue and discomfort that can be experienced with aging. Anaerobic
25 stress can also result from prolonged submaximal isometric exercise when the local circulation is partially or totally occluded by the increase in intra-muscular pressure (e.g., during rock climbing, free diving, or synchronized swimming). Excessive lactate production can result in the
30 acidification of the intracellular environment.

 Creatine (i.e., N-(aminoiminomethyl)-N-glycine, N-amidinosarcosine, N-methyl-N-guanylglycine, or methylglycocyamine) is found in large amounts in skeletal muscle and other "excitable" tissues (e.g., smooth muscle,
35 cardiac muscle, or spermatozoa) characterized by a capacity

for a high and variable energy demand. Creatine is converted into phosphorylcreatine in energy-generating biochemical pathways within cells. In mammalian skeletal muscle, the typical combined content of creatine (i.e.,
5 creatine and phosphorylcreatine) may vary from less than 25 to about 50 mmol per kilogram fresh muscle (i.e., 3.2 to 6.5 grams per kilogram fresh muscle).

Creatine formed is formed in the liver and taken up into tissues, such as muscle, by means of an active
10 transport system. Creatine synthesis in the body may also be augmented by the ingestion of creatine present in meat (e.g., 5-10 milligrams per kilogram body weight per day in the average meat-eating human and approximately zero in a vegetarian diet).

15 During sustained intensive exercise, or exercise sustained under conditions of local hypoxia, the accumulation of hydronium ions formed during glycolysis and the accumulation of lactate (anaerobic metabolism) can severely reduce the intracellular pH. The reduced pH can
20 compromise the function of the creatine-phosphorylcreatine system. The decline in intracellular pH can affect other functions within the cells, such as the function of the contractile proteins in muscle fibers.

Dipeptides of beta-alanine and histidine, and their
25 methylated analogues, include carnosine (beta-alanyl-L-histidine), anserine (beta-alanyl-L-1-methylhistidine), or balenine (beta-alanyl-L-3-methylhistidine). The dipeptides are present in the muscles of humans and other vertebrates. Carnosine is found in appreciable amounts in muscle of, for
30 example, humans and equines. Anserine and carnosine are found in muscle of, for example, canines, camelids and numerous avian species. Anserine is the predominant beta-alanylhistidine dipeptide in many fish. Balenine is the

predominant beta-alanylhistidine dipeptide in some species of aquatic mammals and reptiles. In humans, equines, and camelids, the highest concentrations of the beta-alanylhistidine dipeptides are found in fast-contracting glycolytic muscle fibers (type IIA and IIB) which are used extensively during intense exercise. Lower concentrations are found in oxidative slow-contracting muscle fibers (type I). See, e.g., Dunnett, M. & Harris, R.C. Equine Vet. J., Suppl. 18, 214-217 (1995). It has been estimated that carnosine contributes to hydronium ion buffering capacity in different muscle fiber types; up to 50% of the total in equine type II fibers.

Summary of the Invention

In general, the invention features methods and compositions for increasing the anaerobic working capacity of muscle and other tissues. The method includes simultaneous accumulation of creatine and beta-alanylhistidine dipeptides, or beta-alanine and L-histidine analogues, within a tissue in the body. The methods include ingesting or infusing compositions into the body. The compositions are mixtures of compounds capable of increasing the availability and uptake of creatine and of precursors for the synthesis and accumulation of beta-alanylhistidine dipeptides, in human and animal muscle. The composition induces the synthesis and accumulation of beta-alanylhistidine dipeptides in a human or animal body when introduced into the body.

The compositions include mixtures of creatine and beta-alanine, creatine, beta-alanine and L-histidine, or creatine and active derivatives of beta-alanine or L-histidine. Each of the beta-alanine or L-histidine can be the individual amino acids, or components of dipeptides,

oligopeptides, or polypeptides. The beta-alanine or L-histidine can be active derivatives. An active derivative is a compound derived from, or a precursor of, the substance that performs in the same or similar way in the body as the
5 substance, or which is processed into the substance and placed into the body. Examples include, for example, esters and amides.

In one aspect, the invention features a method of regulating hydronium ion concentrations in a tissue. The
10 method includes the steps of providing an amount of beta-alanine to blood or blood plasma effective to increase beta-alanylhistidine dipeptide synthesis in a tissue, and exposing the tissue to the blood or blood plasma, whereby the concentration of beta-alanylhistidine is increased in
15 the tissue. The method can include the step of providing an amount of L-histidine to the blood or blood plasma effective to increase beta-alanylhistidine dipeptide synthesis.

In another aspect, the invention features a method of increasing the anaerobic working capacity of a tissue.
20 The method includes the steps of providing an amount of beta-alanine to blood or blood plasma effective to increase beta-alanylhistidine dipeptide synthesis in a tissue, providing an amount of L-histidine to the blood or blood plasma effective to increase beta-alanylhistidine dipeptide
25 synthesis in a tissue, and exposing the tissue to the blood or blood plasma. The concentration of beta-alanylhistidine is increased in the tissue.

In embodiments, the methods can include the step of increasing a concentration of creatine in the tissue. The
30 increasing step can include providing an amount of creatine to the blood or blood plasma effective to increase the concentration of creatine in the tissue (e.g., by providing the amount of creatine to the blood or blood plasma).

The providing steps of the methods can include ingestion or infusion (e.g., injection) of a composition including the amount of beta-alanine, or the amounts of beta-alanine and L-histidine, or a combination of ingestion
5 and infusion.

The methods can include increasing a concentration of insulin in the blood or blood plasma. The concentration of insulin can be increased, for example, by injection of insulin.

10 The tissue can be a skeletal muscle.

In another aspect, the invention features a composition consisting essentially of a peptide source including beta-alanine, between about 39 and about 99 percent by weight of a carbohydrate, and up to about 60
15 percent by weight of water. The composition includes between about 1 and about 20 percent by weight of the beta-alanine. The peptide source can include L-histidine. The composition can include between about 1 and about 20 percent by weight of L-histidine.

20 The carbohydrate can be a simple carbohydrate (e.g., glucose). In another aspect, the invention features a composition consisting essentially of a peptide source including beta-alanine, between about 1 and about 98 percent by weight of a creatine source, and up to about 97
25 percent by weight of water. The composition includes between about 1 and about 98 percent by weight of the beta-alanine. The peptide source can include L-histidine and the composition includes between about 1 and about 98 percent by weight of L-histidine.

30 The peptide source can be a mixture of amino acids, dipeptides, oligopeptides, polypeptides, or active derivatives thereof.

The composition can be a dietary supplement. The creatine source can be creatine monohydrate.

5 The concentrations of components in blood or blood plasma can be increased by infusion (i.e., injection) or ingestion of an agent operable to cause an increase in the blood plasma concentration. The composition can be ingested in doses of between about 10 grams and about 800 grams per day. The doses can be administered in one part or multiple parts each day.

10 An increase in the muscle content of creatine and beta-alanylhistidine dipeptides can increase the tolerance of the cells to the increase in hydronium ion production with anaerobic work, and to lead to an increase in the duration of the exercise before the onset of fatigue. The
15 compositions and methods can contribute to correcting the loss of beta-alanine, L-histidine, or creatine due to degradation or leaching of these constituents during cooking or processing. The compositions and methods can also contribute to correcting the absence of these components
20 from a vegetarian diet.

The methods and compositions can be used to increase beta-alanylhistidine dipeptide by, for example, sportsmen, athletes, body-builders, synchronized swimmers, soldiers, elderly people, horses in competition, working and racing
25 dogs, and game birds, to avoid or delay the onset of muscular fatigue.

Other advantages and features of the invention will be apparent from the detailed description, and from the claims.

30 Brief Description of the Drawing

Fig. 1 is a graph depicting changes in the concentrations of beta-alanine in blood plasma of five

horses, before and at 2 hour intervals following the feeding of beta-alanine and L-histidine (100 milligrams per kilogram body weight and 12.5 milligrams per kilogram body weight, respectively, three times per day) over a period of 30 days.

5 Fig. 2 is a graph depicting changes in the concentrations of L-histidine in blood plasma of five horses, before and at 2 hour intervals following the feeding of beta-alanine and L-histidine (100 milligrams per kilogram body weight and 12.5 milligrams per kilogram body weight, respectively, three times per day) over a period of 30 days.

10 Figs. 3a and 3b are graphs depicting the contrast in the changes in the concentrations of beta-alanine in blood plasma of six horses, before and at hourly intervals following the feeding of beta-alanine and L-histidine (100 milligrams per kilogram body weight and 12.5 milligrams per kilogram body weight, respectively, three times per day) on the first and last day of a 30 day period of dietary supplementation.

15 Figs. 4a and 4b are graphs depicting the contrast in the changes in the concentrations of L-histidine in blood plasma of six horses, before and at hourly intervals following the feeding of beta-alanine and L-histidine (100 milligrams per kilogram body weight and 12.5 milligrams per kilogram body weight, respectively, three times per day) on the first and last day of a 30 day period of dietary supplementation.

20 Fig. 5 is a graph depicting the contrast in the changes in the mean concentrations of beta-alanine in equine blood plasma (n = 6), before and at hourly intervals following the feeding of beta-alanine and L-histidine (100 milligrams per kilogram body weight and 12.5 milligrams per kilogram body weight, respectively, three times per day) on

the first and last day of a 30 day period of dietary supplementation.

Fig. 6 is a graph depicting the contrast in the changes in the mean concentrations of L-histidine in equine blood plasma ($n = 6$), before and at hourly intervals following the feeding of beta-alanine and L-histidine (100 milligrams per kilogram body weight and 12.5 milligrams per kilogram body weight, respectively, three times per day) on the first and last day of a 30 day period of dietary supplementation.

Fig. 7 is a graph depicting the correlation between the increase in 6 thoroughbred horses in the carnosine concentration in type II skeletal muscle fibers (the average of the sum of type IIA and IIB fibres) and the increase, between the 1st and 30th day of supplementation, in the area under the blood plasma beta-alanine concentration-time curve over the first 12 hours of the day ($AUC_{(0-12 \text{ hr})}$).

Fig. 8 is graph depicting the mean results of the administration of beta-alanine, broth, or carnosine to test subjects.

Fig. 9 is a graph depicting mean changes in plasma beta-alanine over nine hours of treatment.

Fig. 10 is a graph depicting the mean changes in plasma beta-alanine over 9 hours following the oral ingestion of 10 milligrams per kilogram body weight of beta-alanine.

Fig. 11 is a graph depicting the mean ($n = 6$) plasma beta-alanine concentration over the 24 hour of Day 1 and Day 30 of the treatment period.

Description of the Preferred Embodiments

Beta-alanylhistidine dipeptides such as carnosine, anserine, and balenine have pKa values between approximately

6.8 and 7.1 and are involved in the regulation of intra-cellular pH homeostasis during muscle contraction and the development of fatigue. The content of other substances involved in hydronium ion buffering, such as amino acid residues in proteins, inorganic and organic phosphates and bicarbonate, is constrained by their involvement in other cell functions. The beta-alanylhistidine dipeptides can provide an effective way of accumulating pH-sensitive histidine residues into a cell. Variations in the muscle beta-alanylhistidine dipeptide concentrations affect the anaerobic work capacity of individual athletes.

The beta-alanylhistidine dipeptides are synthesized within the body from beta-alanine and L-histidine. These precursors can be generated within the body or are made available via the diet, including from the breakdown of an ingested beta-alanylhistidine dipeptide. Beta-alanine within the body is transported to tissues such as muscle. In a typical fed state, the concentration of beta-alanine is low in comparison with the concentration of L-histidine in human and equine blood plasma. These concentrations should be viewed in relation to the affinity of the carnosine synthesizing enzyme, carnosine synthetase, for its substrates as determined by the Michaelis-Menton constant (K_m). The K_m for histidine is about $16.8 \mu M$. The K_m for beta-alanine is between about 1000 and $2300 \mu M$. The low affinity of carnosine synthetase for beta-alanine, and the low concentration of beta-alanine in muscle, demonstrate that the concentration of beta-alanine in muscle which is limiting to the synthesis of the beta-alanylhistidine dipeptides.

Increasing the amount of beta-alanylhistidine dipeptides within a muscle favorably affects muscular performance and the amount of work that can be performed by

the muscle. Accordingly, the synthesis and accumulation of beta-alanylhistidine dipeptides is increased in a tissue in a human or animal body.

The synthesis and accumulation of beta-alanylhistidine dipeptides in a human or animal body can be increased with an increase in the content within the body of creatine, by increasing the blood or blood plasma concentrations of beta-alanine, increasing the blood or blood plasma concentrations beta-alanine and creatine, or increasing the blood or blood plasma concentrations beta-alanine, L-histidine, and creatine. The increase in dipeptide can be simultaneous with the increase in beta-alanine concentration.

The blood plasma concentrations of beta-alanine, L-histidine and creatine can be increased by ingestion or infusion of beta-alanine, L-histidine, and creatine, or active derivatives thereof. The composition can be administered orally, enterally, or parenterally. The beta-alanine and creatine, or beta-alanine, L-histidine and creatine, are preferably orally ingested.

The composition can include carbohydrates (e.g., simple carbohydrates), insulin, or agents that stimulate the production of insulin.

The composition can be ingested as a dietary supplement. Preferably, the composition can be administered in one or more doses per day. The beta-alanine dosage can be between about 5 milligrams and about 200 milligrams per kilogram body weight. The creatine (e.g., creatine monohydrate) dosage can be between about 5 milligrams to 200 milligrams per kilogram body weight. The L-histidine dosage can be between about 1 milligrams to 100 milligrams per kilogram body weight. The simple carbohydrate (e.g.,

glucose) dosage can be between about 0.5 and 2.0 grams per kilogram body weight.

In an 80 kilogram person, suitable dosages per day can be between 0.4 grams to 16.0 grams of beta-alanine, 0.4
5 grams to 16.0 grams of creatine monohydrate, 0.08 grams to 8.0 grams of L-histidine, or 40 grams to 160 grams of glucose or other simple carbohydrate. The composition can be in solid form or liquid form or the form of a suspension which is ingested, or in liquid form or suspension for
10 infusion into the body. The composition is ingested in humans in an amount of between 2 grams and 1000 grams per day (e.g., between 10 grams and 800 grams), which may be taken in one or more parts throughout the day. In animals the daily intake will be adjusted for body weight.

15 For humans and animals, the compositions can be:

- (a) 1% to 99% by weight of beta-alanine;
1% to 99% by weight of creatine monohydrate; and
0% to 98% by weight of water;
- (b) 1% to 98% by weight of beta-alanine;
20 1% to 98% by weight of L-histidine;
1% to 98% by weight of creatine monohydrate; and
0% to 97% by weight of water;
- (c) 1% to 20% by weight of beta-alanine;
39% to 99% by weight of glucose or other simple
25 carbohydrate; and
0% to 60% by weight of water; or
- (d) 1% to 20% by weight of beta-alanine;
1% to 20% by weight of L-histidine
39% to 99% by weight of glucose or other simple
30 carbohydrate; and
0% to 60% by weight of water.

The following are specific examples of the methods of methods and compositions for increasing the anaerobic working capacity of muscle and other tissues.

Example 1

5 The effect of supplementation of a normal diet with multiple daily doses of beta-alanine and L-histidine on the carnosine concentration in type I, IIA, and IIB skeletal muscle fibers of thoroughbred horses was assessed. Six
10 experimental thoroughbred horses of normal health (three fillies and three geldings), aged 4 to 9 years, underwent one month (30 days) of dietary conditioning (pre-supplementation period) prior to the commencement of the supplementation period. During the dietary conditioning
15 period each horse was fed a diet comprising 1 kilogram of pelleted feed (Spillers racehorse cubes) and 1 kilogram of soaked sugar beet pulp as a source of complex and simple carbohydrates, three times per day (at 08:30, 12:30, and 16:30, respectively). Soaked hay (3 kilograms dry weight) was also provided twice daily (at 09:00 and 17:00). Water
20 was provided *ad libitum*.

 During the supplementation period an identical feeding regime was implemented. However, each hard feed meal was supplemented with beta-alanine and L-histidine (free base). Beta-alanine and L-histidine were mixed
25 directly into the normal feed. Individual doses of beta-alanine and L-histidine were calculated according to body weight. Beta-alanine was administered at 100 milligrams per kilogram body weight and L-histidine at 12.5 milligrams per kilogram body weight. Dietary supplementation was begun on
30 day 1 of the protocol and discontinued at the end of day 30. Heparinized blood samples (5 milliliters) were collected on days 1, 6, 18, 24, and 30. On day 1 and day 30, blood

samples were collected prior to the first feed and at hourly intervals for a total of 12 hours each day. On the three intervening sampling days, blood was collected prior to the first feed and 2 hours after each subsequent feed. On the day before the start of supplementation (day 0) a muscle biopsy was taken, following application of local anaesthesia of the skin, from the right middle gluteal muscle (m. gluteus medius) of each horse using a Bergstrom-Stille percutaneous biopsy needle. Subsequent muscle biopsies were collected immediately after the end of the supplementation period (day 31) as close as possible to the original sampling site. Clinical monitoring of the horses was performed daily. This comprised a visual examination and measurement of body weight, twice-daily measurement of rectal temperature, and weekly blood sampling for clinical biochemistry and hematology. During the course of the study the horses received no formal training or exercise, although they were allowed one hour of free exercise each day.

Fragments of individual muscle fibers dissected from freeze-dried muscle biopsies were characterized as either type I, IIA or IIB by histochemical staining for myosin ATPase activity at pH 9.6 following pre-incubation at pH 4.50 by a modification of the method described in, Kaiser and Brook, Arch. Neurol., 23:369-379 (1970).

Heparinized blood plasma samples were extracted and analyzed for beta-alanine and L-histidine concentrations by high-performance liquid chromatography (HPLC). Individual weighed muscle fibers were extracted and analyzed for carnosine by HPLC according to the method described in, Dunnett and Harris, "High-performance liquid chromatographic determination of imidazole dipeptides, histidine, 1-methylhistidine and 3-methylhistidine in muscle and

individual muscle fibers," J. Chromatogr. B. Biomed. Appl.,
688:47-55 (1997).

Differences in carnosine concentrations within fiber
types before and after supplementation were established
5 within horses using one-way analysis of variance (ANOVA).
In instances where differences were detected, significance
was determined using a multiple comparison test (Fisher's
PLSD).

No palatability problems were encountered with the
10 addition of beta-alanine and L-histidine to the feed. No
adverse physiological or behavioral effects of the
supplemented diet were observed in any of the horses during
the thirty days of supplementation. No significant changes
in body weight were recorded, and rectal temperatures
15 remained within the normal range. No acute or chronic
changes in clinical biochemistry or hematology were
observed. Beta-alanine was not detected in the plasma of
any of the horses prior to the start of supplementation.
The lower limit of quantitation for beta-alanine in plasma
20 by the assay used was 3 micromolar (μM). Plasma L-histidine
concentrations in the six horses prior to the start of
supplementation were between 36.6 and 54.4 μM .

Individual changes in blood plasma beta-alanine and
L-histidine concentrations for five of the six horses over
25 on all the sampling days are shown in Figs. 1 and 2,
respectively. There was a trend towards an increase in the
pre-feeding concentrations of blood plasma beta-alanine and
L-histidine with increasing time of supplementation.
Furthermore, over the thirty day supplementation period, the
30 blood plasma concentration response to supplementation was
also increased. The response was greater for beta-alanine.

Comparisons of the changes in blood plasma beta-
alanine and L-histidine concentrations prior to the first

feed of the day, and hourly thereafter between the first and last days of the supplementation period, for the six individual horses, are shown in Figs. 3a and 3b, and Figs. 4a and 4b, respectively. The mean (\pm SD) changes ($n = 6$) in blood plasma beta-alanine concentration over time during the 24 hours of the first (day 1) and last (day 30) days of the supplementation period are contrasted in Fig. 5. The area under the mean blood plasma beta-alanine concentration versus time curve over 24 hours ($AUC_{(0-24hr)}$) was much greater on day 30 of the supplementation.

The mean (\pm SD) changes ($n = 6$) in blood plasma L-histidine concentration over time during the 24 hours of the first (day 1) and last (day 30) days of the supplementation period are contrasted in Fig. 6. The area under the mean blood plasma beta-alanine concentration vs. time curve over 24 hours ($AUC_{(0-24hr)}$) was greater on day 30 of the supplementation. The greater AUC for blood plasma beta-alanine on the last day of supplementation (day 30) in contrast to the first day of supplementation (day 1) suggests the increased uptake of beta-alanine from the equine gastro-intestinal tract with progressive supplementation. A similar effect was observed for changes in blood plasma L-histidine concentration during the supplementation period. Peak blood plasma concentrations of beta-alanine and L-histidine occurred approximately one to two hours post-feeding in each case.

A total of 397 individual skeletal muscle fibers (192 pre-supplementation; 205 post-supplementation) from the six horses were dissected and analyzed for carnosine. Mean (\pm SD) carnosine concentration, expressed as millimoles per kilogram dry weight ($mmol\ kg^{-1}\ dw$), in pre- and post-supplementation type I, IIA, and IIB skeletal muscle fibers from the six individual horses are given in Table 1 where n

is the number of individual muscle fibers analyzed. Following thirty days of beta-alanine and L-histidine supplementation the mean carnosine concentration was increased in type IIA and IIB fibers in all six horses. These increases were statistically significant in seven instances. The increase in mean carnosine concentration in type IIB skeletal muscle fibers was statistically significant in five out of six horses. The increase in mean carnosine concentration in type IIA skeletal muscle fibers was statistically significant in two out of six horses.

Table 1

Horse	Day	Type I	n	Type IIA	n	Type IIB	n
6	0 31	32.3 (14.5) -	3	72.1 (47.7) 16.2 (20.9)	11 17	111.8 (22.8) 117.7 (38.7)	14 12
5	0 31	59.5 (3.9) 55.5	2 1	102.6 (12.7) 112.2 (17.1)	12 18	131.2 (26.6) 153.3 (28.0)**	26 22
4	0 31	44.8 (6.6) 37.0 (9.3)	4 2	59.9 (19.5) 88.0 (34.2)*	13 17	108.6 (41.5) 152.4 (65.0)*	19 19
1	0 31	56.7 (5.3) 57.8	2 1	88.5 (20.9) 96.1 (17.3)	15 19	101.3 (15.2) 14.3 (13.3)*	13 11
2	0 31	- 65.9 (13.2)	4	89.6 (16.2) 102.2 (22.1)	13 18	104.2 (22.2) 142.0 (35.4)***	14 12
3	0 31	30.9 (4.0) -	2	85.1 (20.3) 105.0 (17.6)*	6 23	113.5 (20.4) 135.4 (24.9)*	23 9
Mean	0 31	44.8 54.1	13 8	83.0 96.6*	70 112	111.8 135.9**	109 85

- * significantly different to pre-supplementation, $p < 0.05$
 ** significantly different to pre-supplementation, $p < 0.01$
 *** significantly different to pre-supplementation, $p < 0.005$

The absolute (e.g. mmol kg⁻¹ dw) and percentage increases in the mean carnosine concentrations in type IIA and IIB skeletal muscle fibers from the six horses are listed in Table 2.

Table 2

Horse	Type IIA	Type IIA	Type IIB	Type IIB
	Absolute increase	% increase	Absolute increase	% increase
6	4.1	5.7	5.6	5.3
5	9.6	9.4	22.1	16.8
4	28.1	46.9	43.8	40.3
1	7.6	8.6	13.0	12.8
2	12.6	14.1	37.8	36.3
3	19.9	23.4	21.9	19.3
Mean	13.6	18.0	24.1	21.8

10 It was observed that the individual horses which showed
 the greater increase in muscle carnosine concentration
 following thirty days of supplementation also demonstrated the
 greater increase in blood plasma beta-alanine AUC between day 1
 and day 30 of the supplementation period. Referring to Fig. 7,
 15 a significant correlation ($r = 0.986$, $p < 0.005$) for five of
 the six horses was observed between the increase in mean
 carnosine concentration, averaged between type IIA and IIB
 skeletal muscle fibers and the increase, between the 1st and
 30th day of supplementation, in blood plasma beta-alanine AUC,
 20 over the first 12 hours ($AUC_{(0-12 \text{ hr})}$). Only five horses were
 used to calculate the regression line. Horse 6 (filled circle)
 showed no appreciable increase in blood plasma beta-alanine
 concentration greater than that observed on day 1 until the
 last day of supplementation. This was unlike the other five
 25 horses which showed a progressive increase with each sampling
 day. For this reason horse 6 was excluded from the calculation
 of the regression equation.

Increases in muscle carnosine concentration following
 thirty days of supplementation with beta-alanine and L-
 30 histidine will cause a direct increase in total muscle
 buffering capacity. This increase can be calculated by using
 the Henderson-Hasselbach Equation. Calculated values for the
 increases in muscle buffering capacity in type IIA and IIB
 skeletal muscle fibers in the six thoroughbred horses are shown
 35 in Table 3.

Table 3

Horse	Day	Type IIA	Type IIA	Type IIA	Type IIB	Type IIB	Type IIB
		β mcar	β mtotal	$\Delta\beta$ mtotal (%)	β mcar	β mtotal	$\Delta\beta$ mtotal (%)
6	0 31	23.9 25.3	93.9 95.3	+1.5	37.1 39.0	107.1 109.0	+1.8
5	0 31	34.0 37.2	104.0 107.2	+3.1	43.5 50.8	113.5 120.8	+6.4
4	0 31	19.9 29.2	89.9 99.2	+10.3	36.0 50.5	106.0 120.5	+13.7
1	0 31	29.3 31.9	99.3 101.9	+2.6	33.6 37.9	103.6 107.9	+4.2
2	0 31	29.7 33.9	99.7 103.9	+4.2	34.5 47.1	104.5 117.1	+12.1
3	0 31	28.2 34.8	98.2 104.8	+6.7	37.6 44.9	107.6 114.9	+6.8
Mean	0 31	27.5 32.1	97.5 102.1	+4.7	37.1 45.0	107.1 115.0	+7.5

10 Example 2

The effect of supplementation of a normal diet with multiple daily doses of beta-alanine and L-histidine on the carnosine content of type I, IIA, and IIB skeletal muscle fibers of humans was assessed. The plasma concentration of beta-alanine in six normal subjects following the consumption of a broth delivering approximately 40 milligrams per kilogram body weight of beta-alanine was monitored. Doses of 10 and 20 milligrams per kilogram body weight of beta-alanine were also given.

The broth was prepared as follows. Fresh chicken breast (skinned and boned) was finely chopped and boiled for fifteen minutes with water (1 liter for every 1.5 kg of chicken). Residual chicken meat was removed by coarse filtration. The filtrate was flavored by the addition of carrot, onion, celery, salt, pepper, basil, parsley and tomato puree, and reboiled for a further fifteen minutes and then cooled before final filtration through fine muslin at 4°C. The yield from 1.5 kilograms of chicken and one liter of water was 870 mL of broth. A portion of the stock was assayed for the total beta-alanyl-dipeptide content

(e.g., carnosine and anserine) and beta-alanine. Typical analyses were:

total beta-alanyl-dipeptides	74.5 mM
free beta-alanine	5.7 mM

5 The six male test subjects were of normal health and between 25 - 53 years of age, as shown in Table 4. The study commenced after an overnight fast (e.g., a minimum of 12 hours after the ingestion of the last meat containing meal). Subjects were given the option to consume a small quantity of warm water
10 prior to the start of the study. Catheterization was begun at 08:30 and the study started at 09:00.

 As a control, 8 milliliters per kilogram body weight of water was ingested (e.g., 600 mL in a subject weighing 75 kilograms).

15 In one session, 8 milliliters per kilogram body weight of broth containing approximately 40 milligrams per kilogram body weight of beta-alanine (e.g., in the form of anserine and carnosine) was ingested. For a subject weighing 75 kilograms this amounted to the ingestion of 600 milliliters of broth
20 containing 3 grams of beta-alanine. In another session, 3 milliliters per kilogram body weight of a liquid containing the test amount of beta-alanine with an additional 5 milliliters per kilogram body weight of water was ingested. In all sessions, subjects additionally consumed a further 8 milliliters per
25 kilogram body weight of water (in 50 mL portions) during the period of 1 to 2 h after ingestion. A vegetarian pizza was provided after 6 hours. An ordinary diet was followed after 8 hours.

 2.5 milliliter venous blood samples were drawn through an
30 indwelling catheter at 10 minute intervals for the first 90 minutes and then after 120, 180, 240 and 360 minutes. The blood samples were dispensed into tubes containing lithium-heparin as anti-coagulant. The catheter was maintained by flushing with saline. Plasma samples were analyzed by HPLC according to the

method described in Jones & Gilligan (1983) J. Chromatogr.
266:471-482 (1983).

Table 4 summarizes the allocation of treatments during
the beta-alanine absorption study. The estimated equivalent
5 doses of beta-alanine are presented in Table 3.

Table 4

Subject	Age yrs	Weight kg	Broth 40 mg/kg bwt	β -ala 0 mg/kg bwt	β -ala 10 mg/kg bwt	β -ala 20 mg/kg bwt	β -ala 40 mg/kg bwt	Carnosine 20 mg/kg bwt
1	53	76	✓			✓	✓	✓
2	33	60	✓			✓	✓	
3	29	105	✓	✓	✓	✓		
4	31	81	✓	✓	✓		✓	
5	30	94	✓	✓	✓		✓	
6	25	65	✓	✓	✓	✓		

Plasma concentration curves following each treatment are
15 depicted graphically in Fig. 8. Mean results of the
administration of beta-alanine, broth, or carnosine according to
the treatments schedule in Table 4. Plasma beta-alanine was
below the limit of detection in all subjects on the control
treatment. Neither carnosine or anserine were detected in plasma
20 following ingestion of the chicken broth or any of the other
treatments. Ingestion of the broth resulted in a peak
concentration in plasma of 427.9 (SD 161.8) μ M. Administration
of carnosine equivalent to 20 milligrams per kilogram body weight
of beta-alanine in one test subject resulted in an equivalent
25 increase in the plasma beta-alanine concentration.

Administration of all treatments except control resulted
in an increase in the plasma taurine concentration. The changes
in taurine concentration mirrored closely those of beta-alanine.
Administration of broth, a natural food, caused the an equivalent
30 increase in plasma taurine, indicating that such a response is
occurring normally following the ingestion of most meals.

Example 3

The effect of administration of three doses of 10 milligrams per kilogram body weight of beta-alanine per day (i.e., administered in the morning, noon, and at night) for seven days on the plasma concentration profiles of beta-alanine and taurine were investigated. The plasma concentration profiles following administration of 10 milligrams per kilogram body weight of beta-alanine were studied in three subjects at the start and end of a seven day period during which they were given three doses of the beta-alanine per day.

Three male subjects of normal health, aged between 33 - 53 years were studied. Test subjects received three doses per day of 10 milligrams per kilogram body weight of beta-alanine for eight days. In two subjects, this was followed by a further 7 days (days 9 - 15) when three doses of 20 milligrams per kilogram body weight per day were given. Subjects reported at 8 am to the blood collection laboratory on days 1 (prior to any treatment given), 8 and 15 following an overnight fast. Subjects were asked not to consume any meat containing meal during the 12 hours preceeding the study. On each of these three test days subjects were catheterized and an initial blood sample taken when the beta-alanine was administered at or close to 9 am, 12 noon, and 3 pm. Blood samples were drawn after 30, 60, 120 and 180 minutes, and analyzed for changes in the plasma concentration of beta-alanine and taurine. 24 hour urine samples were collected over each day of the study and analyzed by HPLC to determine the excretion of beta-alanine and taurine. The treatments are summarized in Table 5.

Table 5

Treatment Day beta-alanine	Day 1 10 mg/kg bwt	Day 8 10 mg/kg bwt	Day 15 20 mg/kg bwt
1	✓	✓	✓
2	✓	✓	✓
3	✓	✓	

The plasma beta-alanine concentrations are summarized in Fig. 9. Each dose resulted in a peak beta-alanine concentration at one-half hour or one hour after ingestion followed by a decline to a 0-10 micromolar basal level at three hours, just prior to administration of the next dose. The response on day 8 of the treatment tended to be less than on day 1 as indicated by the area under the plasma concentration curve.

Example 4

The effect of administration of three doses of 40 milligrams per kilogram body weight of beta-alanine per day (i.e., administered in the morning, noon, and at night) for 2 weeks on the carnosine content of muscle, and isometric endurance at 66% of maximal voluntary contraction force.

Six normal male subjects, aged 25 to 32 years, that did not have evidence of metabolic or muscle disease were recruited into the study. The subjects were questioned regarding their recent dietary and supplementary habits. None of subjects was currently taking supplements containing creatine, or had done so in recent testing supplementation procedures. The physical characteristics of the test subjects are summarized in Table 6.

Table 6

Subject	Age (years)	Weight (kg)
1	29	78
2	31	94
3	29	105
4	25	65
5	31	81
6	25	75
7	53	76

Two days before treatment, a preliminary determination of maximal voluntary (isometric) contraction force (MVC) of knee extensors with the subject in the sitting position was carried out. MVC was determined using a Macflex system with subjects motivated by an instantaneous visual display of the force output. For each subject, two trials were carried out to determine endurance at 66% MVC sustained until the target force could no longer be maintained despite vocal encouragement. This first contraction was subsequently followed by a rest period of 60 seconds, with the subject remaining in the isometric chair. After the rest period, a second contraction was sustained to fatigue. Following a second rest of 60 seconds, a third contraction to fatigue was undertaken.

One day before treatment, the subjects reported to the isometric test laboratory between 8 and 10 am. MVC was determined and endurance at 66% MVC over three contractions with 60 second rest intervals, as described above, was determined. Measurements were determined using the subject's dominant leg. A biopsy of the lateral portion of the vastus lateralis was taken again from the dominant leg.

On day 1 of the treatment study, subjects reported to the blood sampling laboratory at 8 am following an overnight fast and a minimum of 12 hours since the last meat containing meal. Following catheterization and a basal blood sample, each subject

followed the supplementation and blood sampling protocol described in Example 3. A dose of 10 milligrams per kilogram body weight of beta-alanine was administered at time 0 (9 am), 3 hours, and 6 hours.

5 On days 2-15, subjects continued to take three doses of 10 milligrams per kilogram body weight of beta-alanine.

 In the morning of day 14, post-treatment isometric exercise tests were conducted on the dominant leg to determine MVC and endurance at 66% MVC relative to the 66% MVC measured on
10 the day prior to treatment. In the afternoon, a muscle biopsy was taken of the vastus lateralis from close to the site of the biopsy taken on the day before treatment.

 On day 15, the procedures followed on day 1 were repeated to determine any overall shift in the plasma concentration
15 profile of beta-alanine and taurine over the 15 days of supplementation. Mean changes in plasma beta-alanine over 9 hours following the oral ingestion of 10 milligrams per kilogram body weight of beta-alanine at 0, 3 and 6 hours on days 1 and 15 when dosing at 3 x 10 milligrams per kilogram body weight per day
20 are shown in Fig. 10.

 One additional test subject (number 7) followed the study, taking three doses 10 milligrams per kilogram body weight for 7 days followed by three doses of 20 milligrams per kilogram body weight for 7 days. No muscle biopsies were taken from this
25 test subject.

 There was no apparent change in the muscle carnosine content in the muscle of the six subjects biopsied. Changes in plasma taurine concentrations in the six subjects mirrored those of beta-alanine, as noted in Example 2.

30 Values from the MVC and endurance at 66% MVC measurements one day before treatment and after 14 days after treatment with three doses of 10 milligrams per kilogram body weight of beta-alanine are listed in Table 7. The mean endurance time at 66% MVC increased in 5 of the 6 subjects. An increase was also seen
35 in subject 7 taking the higher dose.

Table 7

Subject	MVC 1st try N	MVC 2nd try N	time @ 66% MVC 1st seconds	time @ 66% MVC 2nd seconds	time @ 66% MVC 3rd seconds	Total Contraction Time seconds
Pre						
1	784.5	821.9	48.53	29.03	23.78	100.83
2	814.4	886.2	48.40	26.03	16.90	91.33
3	984.9	970.4	38.15	26.03	16.78	80.95
4	714.6	740.4	89.03	56.15	45.65	190.83
5	1204.8	1217.2	37.65	27.64	21.53	86.83
6	722.4	716.8	46.78	29.40	21.90	98.08
Pre mean	870.9	892.1	51.4	32.4	24.3	108.1
Pre SD	190.6	184.6	19.1	11.7	10.8	41.2
Post						
1	895.6	908.0	47.08	30.38	24.03	101.48
2	832.2	908.0	46.65	31.28	18.40	96.33
3	973.7	952.2	42.65	25.03	16.03	83.70
4	814.1	863.9	114.40	64.28	48.53	227.20
5	1246.6	1233.0	42.03	22.78	19.40	84.20
6	760.8	773.3	52.28	31.53	25.95	109.73
Post mean	920.5	939.7	57.5	34.2	25.4	117.1
Post SD	175.7	156.0	28.1	15.2	11.9	54.9
Subject 7						
Pre	858.18	861.54	54.0			
Post	792.54	851.41	62.0			

Other embodiments are within the claims.

What is claimed is:

Claims

1. A method of regulating hydronium ion concentrations in a tissue comprising:
providing an amount of beta-alanine to blood or blood plasma effective to increase beta-alanylhistidine dipeptide synthesis in a tissue; and
exposing the tissue to the blood or blood plasma, whereby the concentration of beta-alanylhistidine is increased in the tissue.
2. The method of claim 1, further comprising providing an amount of L-histidine to the blood or blood plasma effective to increase beta-alanylhistidine dipeptide synthesis in a tissue.
3. The method of claim 1, further comprising increasing a concentration of creatine in the tissue.
4. The method of claim 3, further comprising providing an amount of L-histidine to the blood or blood plasma effective to increase beta-alanylhistidine dipeptide synthesis in a tissue.
5. The method of claim 3, wherein increasing the amount of creatine in the tissue includes providing an amount of creatine to the blood or blood plasma effective to increase the concentration of creatine in the tissue.
6. The method of claim 5, further comprising providing an amount of L-histidine to the blood or blood plasma effective to increase beta-alanylhistidine dipeptide synthesis in a tissue.
7. The method of claim 1, wherein the providing step includes ingestion of a composition including the amount of beta-alanine.

8. The method of claim 1, wherein the providing step includes infusion of a composition including the amount of beta-alanine.

9. The method of claim 1, further comprising increasing a concentration of insulin in the blood or blood plasma.

10. The method of claim 9, further comprising providing an amount of L-histidine to the blood or blood plasma effective to increase beta-alanylhistidine dipeptide synthesis in a tissue.

11. The method of claim 1, wherein the tissue is a skeletal muscle.

12. The method of claim 1, wherein the tissue is a human tissue.

13. The method of claim 1, wherein the tissue is an animal tissue.

14. A method of increasing the anaerobic working capacity of a tissue comprising:

providing an amount of beta-alanine to blood or blood plasma effective to increase beta-alanylhistidine dipeptide synthesis in a tissue;

providing an amount of L-histidine to the blood or blood plasma effective to increase beta-alanylhistidine dipeptide synthesis; and

exposing the tissue to the blood or blood plasma, whereby the concentration of beta-alanylhistidine is increased in the tissue.

15. The method of claim 14, further comprising increasing a concentration of creatine in the tissue.

16. The method of claim 14, wherein the providing steps include ingestion of a composition including the amount of beta-alanine and the amount of L-histidine.

17. The method of claim 14, wherein the providing step includes infusion of a composition including the amount of beta-alanine and the amount of L-histidine.

18. The method of claim 14, further comprising increasing a concentration of insulin in the blood or blood plasma.

19. The method of claim 14, wherein the tissue is a skeletal muscle.

20. The method of claim 14, wherein the tissue is a human tissue.

21. The method of claim 14, wherein the tissue is an animal tissue.

22. A composition consisting essentially of:
a peptide source including beta-alanine;
between about 39 and about 99 percent by weight of a carbohydrate; and
up to about 60 percent by weight of water,
wherein the composition includes between about 1 and about 20 percent by weight of the beta-alanine.

23. The composition of claim 22, wherein the peptide source includes L-histidine and the composition includes between about 1 and about 20 percent by weight of L-histidine.

24. A composition consisting essentially of:
a peptide source including beta-alanine;
between about 1 and about 98 percent by weight of a
creatine source; and
up to about 97 percent by weight of water,
wherein the composition includes between about 1 and
about 98 percent by weight of the beta-alanine.

25. The composition of claim 24, wherein the peptide
source includes L-histidine and the composition includes between
about 1 and about 98 percent by weight of L-histidine.

Figure 1

1/13




Figure 2

2/13








Figure 3a (horses 1-3)

3/13




4/13

FIGURE 30 (HORSES 4-6)




5/13

Figure 4a (horses 1-3)




6/13

Figure 4b (horses 4-6)




7/13

Figure 5




8/13

Figure 6




9/13

Figure 7




10/13

Fig. 8




11/13




12/13

Fig. 10



13/13

Fig. 11



INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 97/01117

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A23L1/305 A61K31/195 A23K1/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A23L A61K A23K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	EP 0 449 787 A (SETRA) 2 October 1991 see the whole document ---	1-21 22-25
A	DE 34 24 781 A (K.NAGAI) 17 January 1985 see page 4, line 23-35 see page 11, line 24-32 see page 12, line 1-19; claims ---	1-25
A	WO 90 06102 A (PEPTIDE TECHNOLOGY) 14 June 1990 see page 2, line 21 - page 3, line 14; claims --- -/--	1-25

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

8 January 1998

Date of mailing of the international search report

21.01.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Moer, A

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 97/01117

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE WPI Section Ch, Week 9409 Derwent Publications Ltd., London, GB; Class B05, AN 94-071835 XP002051438 & JP 06 024 976 A (NIPPON STEEL CORP) , 1 February 1994 see abstract</p> <p>---</p>	1-25
A	<p>DATABASE WPI Section Ch, Week 9122 Derwent Publications Ltd., London, GB; Class B04, AN 91-159952 XP002051439 & JP 03 095 125 A (OTSUKA PHARM CO LTD) , 19 April 1991 see abstract</p> <p>---</p>	1-25
A	<p>DATABASE WPI Section Ch, Week 8005 Derwent Publications Ltd., London, GB; Class A23, AN 80-08279C XP002051440 & JP 54 159 393 A (AJINOMOTO KK) , 17 December 1979 see abstract</p> <p>---</p>	1-25
A	<p>DATABASE WPI Section Ch, Week 9219 Derwent Publications Ltd., London, GB; Class B05, AN 92-156230 XP002051441 & JP 04 095 026 A (NIPPON STEEL CORP) , 27 March 1992 see abstract</p> <p>---</p>	1-25
A	<p>DATABASE WPI Section Ch, Week 9221 Derwent Publications Ltd., London, GB; Class B05, AN 92-173167 XP002051442 & JP 04 112 825 A (NIPPON STEEL CORP) , 14 April 1992 see abstract</p> <p>-----</p>	1-25

INTERNATIONAL SEARCH REPORT

Int. application No.
PCT/IB 97/01117

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Remark : Although claims 1 to 21
are directed to a method of treatment
of the human/animal body , the search has been carried out and based on
the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/IB 97/01117

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 449787 A	02-10-91	IT 1240336 B AT 128360 T DE 69113304 D	07-12-93 15-10-95 02-11-95
DE 3424781 A	17-01-85	JP 1645372 C JP 3012045 B JP 60016934 A	28-02-92 19-02-91 28-01-85
WO 9006102 A	14-06-90	AU 4332089 A EP 0436611 A	26-06-90 17-07-91